



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

be investigated, is how and when such seeds of serotinous cones are eventually liberated and made available, and whether not a great many of them at last perish, the cones never opening.—G. ENGELMANN.

*FRAXINUS QUADRANGULATA* has, at least about Allenton, in St. Louis county, Missouri, hermaphrodite flowers. Mr. G. W. Leterman finds it there common on rocky hills where it is a small tree or shrub with blunt angles of the branchlets, and in rich bottom lands, where the tree is large, and the angles of the branchlets sharp and even winged. Leaves are sometimes in threes when the branchlets show six angles. The terminal buds are gray-downy. In both localities the flowers are hermaphrodite. The calyx is practically absent, or indicated only by two obscure knobs or two minute scales, alternating with the stamens; the anthers are sessile and (before opening) reniform, their two cells being united above; stamens somewhat persistent at least to the beginning of May, when the young obovate-oblong fruits, already somewhat twisted (which twist is more marked in the mature fruit), have reached about half their full size. How does the species behave in other parts of the country? The style of *Fraxinus Americana* is very slender—much longer than the ovary; that of *F. viridis* does not much exceed the ovary.—G. ENGELMANN.

*NOTULÆ EXIGUÆ.*—THREE-FLOWERED BLOODROOT.—Among the anomalies occasionally met with, the most unexpected is a scape of *Sanguinaria Canadensis*, found by Mr. E. N. Wheeler, in the vicinity of Boston, bearing a pair of opposite bracts about half an inch below the terminal flower, each bract with a well-formed flower in its axil!

*TRILLIUM SESSILE*, as we learn from Mr. Lehman, of Salem, North Carolina, and from a specimen sent by him, abundantly occurs in the neighborhood of Kingston, Tenn., with bright yellow petals; and I have recently heard of this form from other western sources. Specimens, and especially living roots, taken up in autumn, are desired.

*PERULARIA VIRESCENS* is the proper name for *Habenaria* (*Perularia*) *virescens* of Gray's Manual. The examination of fresh specimens shows the "*cuculli bivalves*" of Lindley, the two lips of the base of the anther-cell which fairly cover the gland. In Florida specimens just received from Miss Reynolds, of Florida, the outer lip is the larger, or the one which principally protects the gland. How is it in the northern plant? It has long since been announced by me in Am. Jour. Sci., that *Orchis rotundifolia*, Pursh, is a true *Orchis*.—A. GRAY.

*PLATANThERA BRACTEATA*, Torr.—This is usually regarded as a summer flowering species. In my garden, where it has bloomed for the first time this season, it is the earliest of many that I have. The first flowers were open on the 26th of April, and half of the spike had opened by the 1st of May. Hitherto *Orchis spectabilis* has been the